

*Sandy, Final Report -
Phone file
and
close out file -
Thanks
WOK*

FINAL TECHNICAL REPORT
AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

Grant Number: F49620-96-1-0241

Grant Title: Study Of Short-Pulsed Laser Retinal Injury Mechanisms By
Time-Resolved Imaging Of Photomechanical Transients In RPE

Principal Investigator: Charles P. Lin, PhD

Institution: Wellman Laboratories of Photomedicine
Massachusetts General Hospital
BHX 630
Boston, MA 02114

Funding period: 01 JUL 96 - 30 JUN 99

20000718 090

DEC QUALITY INSPECTION

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-00-

88

Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Room 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project, Washington, DC 20503.

g data sources,
r aspect of this
1215 Jefferson
20503.

0252

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 14 May 2000	3. REPORT TYPE AND DATES COVERED FINAL - 01 Jul 96 - 30 Jun 99	
4. TITLE AND SUBTITLE Study of Short-Pulsed Laser Retinal Injury Mechanisms By Time-Resolved Imaging of Photomechanical Transients in RPE		5. FUNDING NUMBERS F49620-96-1-0241	
6. AUTHOR(S) Charles P. Lin, PhD			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Wellman Laboratories of Photomedicine Massachusetts General Hospital BHX 630 Boston MA 02114		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL 801 N Randolph St., Rm 732 Arlington VA 22203-1977		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release: Distribution Unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We studied RPE cell damage mechanism for laser duration from 100 fsec to 5 usec, and we have investigated the dependence of threshold fluence for cell damage on the laser spot size on the RPE. These results will be summarized in the report. Our current method for studying cell damage mechanism in the RPE is stroboscopic time resolved microscopy, which allows us to image the transient microscopic bubble formation inside the RPE tissue ex vivo where the cells following pulse laser irradiation. This technique works well in RPE tissue ex vivo where the strobe light transilluminates the specimen. Since our ultimate goal is to detect the existence of these micobubbles in vivo, we have set up a stroboscopic imaging system in epi-illumination geometry. However, the image quality obtained in this way is inferior to that obtained by transillumination, most likely because of the strong angular dependence of the back-scattered light by the spherical bubbles. It is thus difficult to identify the extent of bubble formation at the RPE by this method.			
14. SUBJECT TERMS Cell damage, Stroboscopic, Irradiation		15. NUMBER OF PAGES 8	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASS	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASS	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASS	20. LIMITATION OF ABSTRACT

DTIC QUALITY INSPECTED 4

Standard Form 298 (Rev. 2-89) (EG)
Prescribed by ANSI Std. Z39.18
Designed using Perform Pro, WHS/DIOR, Oct 94

Objectives

1. Study laser-induced cavitation bubble formation in RPE cells in vitro using the technique of nsec/psec time-resolved microscopy. Determine the threshold fluence for intracellular cavitation using nanosecond and picosecond laser pulses at 1064 & 532 nm. *(year 1)*
2. Integrate fluorescence imaging capability into existing time-resolved imaging apparatus. Perform RPE cell viability assays and examine the relationship between cell injury and the onset of intracellular cavitation. *(year 1)*
3. Investigate stress wave generation and propagation from target RPE to surrounding cells. Assess the effect of stress transients on neighboring cells not subject to direct laser irradiation and cavitation. *(years 1 and 2)*
4. Investigate the effects of laser spot size on RPE cavitation and cell injury thresholds. *(years 2)*
5. Investigate the possibility of imaging transient bubble formation in RPE in vivo. This will facilitate a direct comparison of in vivo retinal injury threshold with the onset of cavitation. In vivo experiment will be carried out at the Armstrong Laboratory, Brooks AFB. *(year 3)*
6. Investigate the mechanism of multiple pulse laser injury in the RPE. Examine multiple pulse RPE damage using fluences above and below single-pulse mechanical injury threshold to assess the additivity of photomechanical effects. *(year 3)*
7. Investigate femtosecond laser-induced mechanical effects using similar procedures as described above. Femtosecond experiments will be performed using laser sources at Brooks AFB (with the ultrafast laser team in the Armstrong Laboratory). *(year 2)*

Status of Effort

We have studied RPE cell damage mechanism for laser pulse durations from 100 fsec to 5 μ sec, and we have investigated the dependence of threshold fluence for cell damage on the laser spot size on the RPE. These results will be summarized in the section below.

Our current method for studying cell damage mechanism in the RPE is stroboscopic time resolved microscopy, which allows us to image the transient microscopic bubble formation inside the RPE cells following pulse laser irradiation. This technique works well in RPE tissue ex vivo where the strobe light transilluminates the specimen. Since our ultimate goal is to detect the existence of these micobubbles in vivo, we have set up a stroboscopic imaging system in epi-illumination geometry. However, the image quality obtained in this way is inferior to that obtained by transillumination, most likely because of the strong angular dependence of the back-scattered light by the spherical bubbles. It is thus difficult to identify the extent of bubble formation at the RPE by this method.

We are currently investigating two other methods to detect bubble formation in vivo - one based on probe beam reflection and one based on photoacoustic detection. We have shown that the probe beam technique is able to detect bubble formation in explant RPE cells very close to threshold. The method is ready to be applied in vivo.

Accomplishments/New Findings

Pulse width dependence studies

The threshold for laser-induced cavitation and cell death in the RPE was determined using a stroboscopic time-resolved microscope developed for this project. Laser pulse durations of 20 nsec and 30 psec (at 532 nm) were used in the initial experiments and the cell damage threshold was found to be the same as bubble formation threshold. A similar microscope system was assembled at Brooks AFB and the measurements were extended down to 100 fsec. Another system was assembled in the Medical Laser Center in Luebeck and used to probe longer pulse durations (nsec to μ sec). The dependence of RPE cell damage threshold as a function of pulse duration over this entire range (from 10^{-13} to 5×10^{-6} sec) is plotted in Figure 1. The threshold is nearly constant below 100 nsec. The increase in threshold fluence above 100 nsec can be attributed to heat diffusion out of the absorbing melanosome particles.




Figure 1. Dependence of RPE cell damage threshold as a function of laser pulse duration. The three data points below 10^{-11} sec were taken with a laser wavelength of 580 nm. The rest of the data points were taken with 527/532 nm.

Spot size dependence studies

The motivation for the study is the long-standing puzzle from observations made *in vivo*, that the MVL threshold increases as the retina spot size decreases. For long pulse durations, this effect can be explained by heat diffusion away from the irradiated spot, a process which is more efficient for smaller spot sizes, which in turn requires a higher fluence to cause damage. Surprisingly, the same effect was also observed for short pulse durations where thermal diffusion is not expected to play a significant role, and no satisfactory explanation has been put forth for the observation. At the urging of Dr. David Sliney, we investigated the dependence of threshold for cavitation and cell lethality as a function of laser spot size in the *ex vivo* model. We expected to find no spot size dependence because the cavitation mechanism requires only a threshold fluence which is independent of spot size. Our results are shown in Figure 2, together with the *in vivo* data replotted from Beatrice et al. As expected, the *ex vivo* results show a constant fluence for laser spot sizes ranging from 200 μm to 20 μm (data taken with 100 psec laser pulses at 532 nm). The interesting observation is the absolute laser fluences for producing damage. The *in vivo* and *ex vivo* fluence values agree for large spot sizes but diverge significantly for small spot sizes. This raises the question whether a lower fluence exposure *in vivo* indeed creates damage at the RPE which is not observable ophthalmoscopically because the spot size (and consequently the lesion) is

too small. This question can be answered by probing bubble formation directly in the eye in vivo using the probe beam setup described in the next section.




Figure 3. Open circles: ED₅₀ threshold for ex vivo RPE cell damage (100 psec, 532 nm).

Filled squares: MVL threshold in vivo, from Zuclich et al.

Filled triangles: MVL threshold in vivo, from Beatrice et al.

Bubble detection by probe beam reflectance

A pump-probe system was set up to detect transient light scattering from the cavitation bubbles, using a diode laser source, confocal detection geometry, and an avalanche photodiode. The diode laser source (830 nm output) is chosen to minimize heating of the RPE cells by the probe beam itself (melanosome absorption decreases toward long wavelengths). In addition, the power of the diode laser is modulated such that the output is effectively turned off during the time between laser pulses in a multiple pulse train. The confocal detection geometry is used to spatially filter out back reflectance and scattering from the optical system and from tissue layers other than the RPE. The use of avalanche photodiode ensures high sensitivity, to detect microbubbles within a single RPE cell, and fast response time, to capture signal from threshold bubbles during their brief lifetime of approximately 100 nsec. Initial results using this system indicates that we can detect near threshold bubbles from a single RPE cell as well as a single melanosome. Figure 3 shows an example of the probe beam reflectance signal. The diode laser was switched on at -10 μ sec and switched off at +8.5 μ sec to minimize sample heating. The Nd:YAG laser was fired at time zero, which causes a detectable increase in probe beam back scattering (upward signal between 0 and 2

μsec) from cavitation bubble induced in a single RPE cell. This system is being developed for probing transient bubble formation in vivo and for studying multiple pulse laser-tissue interactions.




Figure 3. An example of the probe beam reflectance signal from transient cavitation within the RPE. The diode laser was switched on at $-10 \mu\text{sec}$ and switched off at $+8.5 \mu\text{sec}$ to minimize sample heating. The Nd:YAG laser was fired at time zero. Note the upward signal from increased bubble back scattering (between 0 and 2 μsec) which is opposite to the decreased probe transmission signal shown in figure 2.

Personnel Supported

Charles P. Lin	P.I.	25% effort
Jan Roegerer	Student	80% effort (10/98-6/99)

Publications

1. Lin CP, Kelly MW, Sibayan SA, Latina MA, Anderson RR. Selective Cell Killing by Microparticle Absorption of Pulsed Laser Radiation. IEEE J Select Topics Quant Electron., 5:963-968 (1999).
2. Brinkmann R, Hüttmann G, Rögener J, Lin CP, Roider J, Birngruber R. Origin of RPE-Cell Damage by pulsed laser irradiance in the ns to μ s time regime. Submitted to Lasers Surg Med.
3. Leszczynski D, Pitsillides CM, Anderson RR, Lin CP. Induction of apoptosis and necrosis following pulsed laser irradiation of intracellular pigment microparticles. Submitted to Lasers Surg Med.

Proceedings

1. Hüttmann G, Brinkmann R, Rant B, Birngruber R, Rögener J, Lin CP. "Laser-generated microeffects: microcavitation and absorption properties of melanosomes," in Biomedical Optics: New concepts in therapeutic laser applications, OSA Technical Digest (Optical Society of America, Washington DC, 1999) pp 136-138.
2. Leszczynski D, Pitsillides CM, Anderson RR, Lin CP, "Induction of apoptosis and necrosis following pulsed laser irradiation of intracellular pigment microparticles," in Biomedical Optics: New concepts in therapeutic laser applications, OSA Technical Digest (Optical Society of America, Washington DC, 1999) pp 139-141.
3. Roegerer J, Lin CP, "Photomechanical effects - experimental studies of pigment granule absorption, cavitation, and cell damage", Proc Boulder Damage Meeting (1999).

Interactions/Transitions

Results of our RPE studies were presented in three national and international meetings in 1999.

- Optical Society of America topical meeting, Munich, June 1999.
- Boulder Damage Meeting, Boulder, CO, Oct 1999.
- IEEE/LEOS Annual Meeting, San Francisco, CA, Nov 1999.

New discoveries, inventions, or patent disclosures

None.

Honors/Awards

None.